

"Lagevrio" Administration Manual

Product Name	Lagevrio Capsule
Ingredients/Contents	(Active Ingredient) Molnupiravir 200mg (per capsule)
Description	Orange capsule
Direction/ Dose	- Take 4 capsules, 2 doses a day for a total of 10 doses over 5 days. - Begin treatment within 5 days from the day you experience symptoms after being confirmed of COVID-19. - If you are hospitalized during treatment, continuation of administration will be determined by the medical staff.
Effects/ Benefits	Treatment of COVID-19 confirmed individuals with mild and moderate symptoms who have a high risk of developing severe symptoms. However, only administer on individuals who are unable to use other COVID-19 treatments or those who are not clinically appropriate for other COVID-19 treatments.
Storage	Store at room temperature (15°C-30°C)

♣ How should I take Lagevrio

**Take all 4 capsules together, 2 doses per day for 5 days ->
A total of 10 doses**



- It can be taken regardless of meals, and it is important to **complete treatment** of a total of 10 doses over 5 days without interruption .
- Do not chew or cut the capsule**. Please swallow the tablet as a whole.
- The doses must be taken at regular intervals of 12 hours, so make sure to set the first dose at a convenient time according to your lifestyle. Ex) 9:00 AM/9:00 PM

♣ What if I forget to take my medication on time?

① If less than 10 hours passed from the scheduled time to take the medication:	② If more than 10 hours passed from the scheduled time to take the medication:
- Take the medication as soon as you realize the omission, and take the next medication at the originally set time. - Maintain the original medication time afterwards.	- Do not take medication at the time of recognition and take it at the scheduled time for the next dose. - Maintain the original medication time afterwards.
(Do not double the dosage.)	

♣ What are the cautionary items for taking Lagevrio?

1. Hypersensitive reaction, including anaphylaxis, has been reported from individuals administered with this medication. If you experience signs and symptoms of hypersensitive reactions (i.e. palpitations, rashes, shortness of breath) or anaphylaxis after taking this medication, immediately stop administration and notify the your medical institution to begin suitable treatment.
2. This medication **cannot be administered to pregnant women**, and potentially pregnant women must take this medicine after **confirming that they are not pregnant**. If you took this medication **without knowing that you are pregnant**, **immediately stop administration and consult a medical personnel**. Alternatively, you can also contact or consult MSD Korea (02-331-2000).
3. During treatment, **women planning to conceive** should use effective contraception during and for up to **four days after the last dose**, and **men** with a partner (spouse) in childbearing age should use **effective contraception for up to three months**. Furthermore, **breastfeeding mothers** should stop breastfeeding during treatment and **until 4 days after the last dose** of this medication.
4. Taking medication prescribed by a doctor to another person without a doctor's approval is very dangerous. Doing so may be subject to penalization pursuant to the related laws. In principle, if the administration

is stopped due to symptoms such as abnormal reactions after taking the medication, the remaining capsules should be returned and discarded.

♣ **What are the possible side effects?**

		
Diarrhea	Nausea	Dizziness

* In addition to the above symptoms, there may be other side effects. If you experience any side effects after taking this medication, contact your pharmacist or medical institution for consultation. You may also report side effects to the Korea Institute of Drug Safety and

Risk Management (KIDS) through the hotline (1644-6223).

- For more information, visit the official website of Korea Disease Control and Prevention Agency (www.kdac.or.kr) or a local health center.

